






INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 317/00		A2	(11) International Publication Number: WO 99/10321
			(43) International Publication Date: 4 March 1999 (04.03.99)
(21) International Application Number: PCT/US98/17830 (22) International Filing Date: 28 August 1998 (28.08.98) (30) Priority Data: 60/056,272 29 August 1997 (29.08.97) US (71) Applicant (for all designated States except US): THE JOHNS HOPKINS UNIVERSITY [US/US]; 720 Rutland Avenue, Baltimore, MD 21205 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): TOWNSEND, Craig, A. [US/US]; 116 Midhurst Road, Baltimore, MD 21212 (US). DICK, James, D. [US/US]; 15744 Dover Road, Upperco, MD 21155 (US). PASTERNAK, Gary, R. [US/US]; 311 Edgevale Road, Baltimore, MD 21212 (US). KUHAJDA, Francis, P. [US/US]; 1211 Broadway Road, Lutherville, MD 21209 (US). PARRISH, Nicole, M. [US/US]; Apartment 201, 3185 Pine Orchard Lane, Ellicott City, MD 21042 (US). (74) Agents: POSORSKE, Laurence, H. et al.; Baker & Botts, L.L.P., The Warner, 1299 Pennsylvania Avenue, N.W., Washington, DC 20004 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.	

(54) Title: ANTIMICROBIAL COMPOUNDS



(57) Abstract

This invention provides methods for treating a mycobacterial infection by administering to an animal a pharmaceutical composition containing a compound having the formula $R-SO_n-Z-CO-Y$, where R is an alkyl group having 6-20 carbons; Z is a radical selected from $-CH_2-$, $-O-$, and $-NH-$, two of these radicals coupled together, or $-CH_2=CH_2-$; Y is $-NH_2$, $O-CH_2-C_6H_5$, $-CO-CO-O-CH_3$, or $O-CH_3$; and n is 1 or 2. It has been discovered that these compounds inhibit growth of microbial cells which synthesize α -substituted, β -hydroxy fatty acids, particularly corynemycolic acid, nocardic acid, and mycolic acid. These compounds may be used to inhibit growth of mycobacterial cells, such as *Mycobacterium tuberculosis*, drug-resistant *M. tuberculosis*, *M. avium intracellulare*, and *M. leprae*.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Antimicrobial Compounds

BACKGROUND OF THE INVENTION

Field of the Invention

5 This invention relates to the synthesis and *in vivo* application of compounds which have antibiotic activity against microbes that synthesize mycolic acid, including *Mycobacterium* sp., particularly drug resistant *Mycobacterium* strains, and to the use of these compounds to treat any susceptible pathogenic microorganism or parasite.

10 Review of Related Art

 The emergence of multiply drug resistant (MDR) strains of *Mycobacterium tuberculosis* and other atypical mycobacteria which infect immunocompromised patients (e.g., AIDS patients) highlights the need for continued antibiotic development.

15 *Mycobacterium* sp. synthesize a multitude of complex lipids and glycolipids unique to this genus, making these biochemical pathways attractive targets for drug therapy (Bloch, K., "Control mechanisms for fatty acid synthesis in *Mycobacterium smegmatis*," *Adv. Enzymol.* **45**:1-84, 1977; Brennan, P.J., and Nikaido, H., "The envelope of mycobacteria," *Ann. Rev. Biochem.* **64**:29-63, 1995).

20 The β -ketoacyl synthase (KS) of particulate Type II fatty acid synthases or the corresponding domain of the polyfunctional Type I fatty acid synthases catalyzes the critical two-carbon homologation during buildup of the growing fatty acid chain. This process typically gives acids of length C_{16} to C_{18} . In chain elongation of normal fatty acids, carried out for example by mycobacteria, CoA and/or acyl-carrier protein (ACP)

25 thioesters of these acids are further reacted with malonyl-CoA to greatly extend their length to 60-90 carbons. These high molecular weight acids are known collectively as mycolic acids.

 Mycolic acids are a group of complex, long, branched chain fatty acids that are vital for the growth and survival of mycobacteria. Mycolic acids comprise the

30 single largest component of the mycobacterial cell envelope. Little is known about the

nature of the biosynthetic enzymes involved, but evidence suggests some similarity to conventional fatty acid synthases (Bloch, 1977; Brennan, et al., 1995). These unusually long lipid molecules form a waxy coat of limited permeability.

The presence in mycobacteria of particular modified fatty acids having
5 complex and well-organized structures presents a potentially attractive target for drug design (Young, D.B., and Duncan, K., "Prospects for new interventions in the treatment and prevention of mycobacterial disease," *Ann Rev. Microbiol.* **49**:641-673, 1995). It has been suggested that isoniazid inhibits mycolic acid synthesis as its potential mechanism of action (Takayama, K., Wang, L., and David, H.L., "Effect of isoniazid
10 on the in vivo mycolic acid synthesis, cell growth, and viability of *Mycobacterium tuberculosis*," *Antimicrob. Agents Chemother.*, **2**:29-35, 1972; Takayama, K., Schnoes, H.K., Armstrong, E.I., and Boyle, R.W., "Site of inhibitory action of isoniazid in the synthesis of mycolic acids in *Mycobacterium tuberculosis*," *J. Lipid Res.*, **16**:308-317, 1975; Quemard A., Dessen A., Sugantino M., Jacobs W.R., Sacchettini J.C., Blanchard
15 J.S. "Binding of catalase peroxide-activated isoniazid to wild-type and mutant *Mycobacterium tuberculosis* enoyl-ACP reductases," *J. Am. Chem. Soc.*, **118**:1561-1562, 1996; Baldock C., Rafferty J.B., Sedenikova S.E., Baker P.J., Stuitje A.R., Slabas A.R., Hawkes T.R., Rice D.W. "A mechanism of drug action revealed by structural studies of enoyl reductase," *Science*, **274**:2107-2110, 1996; Quemard A., Sacchettini
20 J.C., Dessen A., Vilcheze C., Bittman R., Jacobs W.R., Blanchard J.S., "Enzymatic characterization of the target for isoniazid in *Mycobacterium tuberculosis*, *Biochemistry*, **34**:8235-8241, 1993; Msluli, K., D.R. Sherman, M.J. Hickey, B.N. Kreiswirth, S. Morris, C.K. Stover, and C.E. Barry, III, "Biochemical and genetic data suggest that InhA is not the primary target for activated isoniazid in *Mycobacterium*
25 *tuberculosis*," *J. Infect. Dis.*, **174**:1085-1090, 1996; Dessen A., A. Quemard, J.S. Blanchard, W.R. Jacobs, and J.C. Sacchettini, "Crystal structure and function of the isoniazid target of *Mycobacterium tuberculosis*," *Science*, **267**:1638-1641, 1995; Banerjee, A., E. Dubnau, A. Quemard, V. Balasubramanian, K.S. Um, T. Wilson, D. Collins, G. deLisle, W.R. Jacobs, Jr., "InhA, a gene encoding a target for isoniazid and

ethionamide in *Mycobacterium tuberculosis*." *Science*, **263**:227-230, 1994). This finding might be expected to stimulate a search for novel compounds that act upon the lipid synthetic pathways of mycobacteria as a fresh approach for antibiotic development. Surprisingly, however, lipid biosynthesis has not been exploited for drug development in these organisms. No drugs which specifically inhibit mycobacterial lipid synthesis have been developed other than isoniazid, and there remains a need for new drugs to treat the growing problem of multi-drug resistant mycobacteria.

SUMMARY OF THE INVENTION

10 This invention is directed to novel compounds having antimicrobial activity, particularly antimicrobial effectiveness against multi-drug resistant mycobacteria.

This invention is also directed to a method for treating mycobacterial infection by drug resistant strains through use of independent therapeutic targets.

15 These and other objects of the invention are achieved by one or more of the following embodiments. In one embodiment, this invention provides a compound having the formula: $R-SO_n-Z-CO-Y$, where R is preferably an alkyl group having 6-20 carbons; Z is preferably a radical selected from $-CH_2-$, $-O-$, and $-NH-$, two of these radicals coupled together or $-CH_2=CH_2-$; Y is preferably $-NH_2$, $-O-CH_2-C_6H_5$, $-CO-CO-O-CH_3$, or $-O-CH_3$; and n is 1 or 2. In particularly preferred embodiments, R is a branched alkyl group, or R is a linear alkyl group interrupted by an aromatic ring.

In another embodiment, this invention provides a method of inhibiting growth of a microbial cell which synthesizes α -substituted, β -hydroxy fatty acids. The method comprising treating the cell with a compound having the formula: $R-SO_n-Z-CO-Y$, as described above. In particular, cells inhibited by the compound of this invention are cells which synthesize α -substituted, β -hydroxy fatty acids selected from the group consisting of corynemycolic acid, nocardic acid, and mycolic acid. Preferably, the method is used to inhibit growth of microbial cells selected from the group consisting of corynebacteria, nocardiae, rhodococcus, and mycobacteria. More

preferably, the method is used to inhibit growth of mycobacterial cells, such as *Mycobacterium tuberculosis*, drug-resistant *M. tuberculosis*, *M. avium intracellulare*, *M. leprae*, or *M. paratuberculosis*.

5 In yet another embodiment, this invention provides a method for treating a mycobacterial infection by administering to an animal a pharmaceutical composition containing a compound having the formula: $R-SO_n-Z-CO-Y$, as described above.

The present inventors have synthesized and tested a number of sulfones and sulfoxides having structures based upon the reaction intermediates of the β -ketoacyl synthase reaction of fatty acid synthase. A number of these compounds have
10 demonstrated *in vitro* activity against virulent *M. tuberculosis* (see Table 1). The desirable characteristics found among the compounds tested included: potency, *in vivo* activity, reproducibility of MIC data, ease of synthesis, and chemical stability. Use of the compounds of this invention in drug therapy against multiply drug resistant tuberculosis will provide a means to treat both patients presently suffering from active
15 disease, and the millions of potential patients who harbor quiescent disease which may become active as a result of immunosuppression or other systemic disease.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a schematic representation of the β -ketoacyl synthase
20 reaction.

Figure 2 shows two-dimensional thin layer chromatography of mycolic acids extracted from *M. avium-intracellulare* (Left Panel) and the same extract following treatment with n-octanesulphonylacetamide (Right Panel).

Figures 3A and 3B show photomicrographs of *M. bovis* BGC with (B) or without (A) prior n-octanesulphonylacetamide treatment.
25


DETAILED DESCRIPTION OF THE EMBODIMENTS

The compounds synthesized according to this invention are based on theoretical transition-state intermediates of the β -ketoacyl synthase reaction of fatty acid

synthase (E.C. 2.3.1.85). The β -ketoacyl synthase reaction is common to fatty acid synthesis in prokaryotes and eukaryotes, including *Mycobacteria*. The family of compounds shown in Table 1 was found to be cytotoxic against a variety of mycobacteria, including drug resistant strains. Instead of inhibiting Type I/Type II fatty acid synthesis in mycobacteria, however, Compound HIII-50 (III-50) inhibits the synthesis of mycolic acids, specialized fatty acids found particularly in *Mycobacteria* sp. One of the key steps in mycolic acid synthesis is fatty acid elongation, which employs carbon-carbon condensation similar to the β -ketoacyl synthase reaction in *de novo* fatty acid synthesis. It is believed that these compounds act at this fatty acid elongation step critical to mycolic acid synthesis.

The transition states for all of the two-carbon elongation reactions are presumed, on existing biochemical evidence and mechanistic grounds, to be quite similar (see Figure 1) and involve acylanion formation by decarboxylation of malonyl-CoA/ACP to provide a nucleophile (Figure 1, step A) to react with the thioester-bound acyl chain to generate tetrahedral intermediate (Figure 1, step B). This transient intermediate is anticipated to collapse (Figure 1, step C) to extend the growing acyl unit (RCO-) by two carbons. Further steps then reduce the β -keto group of the ACP/CoA derivative and finally return it to step A to begin the chain extension cycle again.

Low molecular weight organic compounds which are potential mimics of the tetrahedral intermediate (see stepB) may be expected to be inhibitors of this biochemical process, and such compounds are represented by Formula I.



Formula I

wherein R is hydrocarbon, such as an alkyl group;

n is 1 or 2;

Z is a hydrocarbon linking moiety that may contain a heteroatom; and


Y is a hydrocarbon end group moiety that may contain one or more heteroatoms.

In preferred compounds according to Formula I, R may be an n-alkyl group having 6-20 carbons, preferably 8, 10 or 12 carbons, and may be saturated or unsaturated, branched or unbranched, or the alkyl chain may be interrupted by a

aromatic ring to give ortho-, meta-, or para-disubstitution. Z is preferably $-\text{CH}_2-$, $-\text{CH}_2-\text{CH}_2-$, cis or trans $-\text{CH}=\text{CH}-$, $-\text{NH}-\text{CH}_2-$, $-\text{CH}_2-\text{NH}-$, $-\text{O}-\text{CH}_2-$, or $-\text{CH}_2-\text{O}-$, and Y is preferably $-\text{NH}_2$, $-\text{O}-\text{CH}_2-\text{C}_6\text{H}_5$, $-\text{CO}-\text{O}-\text{CH}_3$, $-\text{CO}-\text{CO}-\text{O}-\text{CH}_3$, or $-\text{O}-\text{CH}_3$. Between R and Z is a sulfur atom in the form of a sulfoxide or sulfone. The sulfone III-50 and close structural analogues with shorter and longer saturated and unsaturated alkyl side chains, as well as alternative mimics of the tetrahedral intermediate of step B are exemplified but not exclusively represented by the structures shown in Table 1.

The compounds of Formula I may be synthesized by a variety of routes including alkylation of a mercaptoester to produce a thioether (see, e.g., R. L. Smith et al., *J. Med. Chem.*, 1977, 20:540-547), followed by oxidation to afford a sulfoxide or sulfone and conversion of the ester to an amide with ammonia or a substituted amine. A suitable synthetic scheme is shown in Scheme 1 and exemplified in Example 1.

Scheme 1



Alternatively, a sulfonate salt, for example the sodium salt, may be reacted with a haloester to give a thioether (i.e. $\text{R}-\text{SO}_2-\text{Na}^+ + \text{X}-(\text{CH}_2)_n-\text{COOR}' \rightarrow \text{R}-\text{SO}_2-(\text{CH}_2)_n-\text{COOR}'$; see, e.g., E. Gipstein, C. G. Willson and H. S. Sachdev, *J. Org. Chem.*, 1980, 45:1486-1489), which can be oxidized and ammonolyzed to the corresponding amide as above. Typically, X = Cl, Br or I. A second alternative would be reaction of a haloamide ($\text{X}-(\text{CH}_2)_n-\text{CONH}_2$) with either a sulfonate salt, as above, or with a thiolate anion followed by oxidation to similarly yield the sulfonyl amide or the sulfoxide amide (see, e.g., S. Huenig and O. Boes, *Liebigs Annalen der Chemie*, 1953, 579:23-26). It will be apparent to the skilled worker that the sense of these reactions can be reversed so that a haloalkane may be reacted with a mercaptoamide or the

salt of a sulfenylamide or sulfenylester to obtain the same products. A third alternative would be reaction of a thiol (R-SH) with propiolate ester or amide to form a sulfanyl-acrylic ester or amide followed by oxidation to the sulfoxide or sulfone. Finally, a halo- or mercaptonitriles can be reacted by the above schemes to give thioethers or
5 sulfonylnitriles, which can be hydrolyzed to the amides. Where other alternative synthetic routes to produce the compounds of Formula I occur to the skilled worker, the products of such synthesis are also within the contemplation of this invention.

The compounds according to Formula I may be used as antibiotics against microbes having in their cell walls α -substituted, β -hydroxy fatty acids, such
10 as corynemycolic acids (e.g., C30), nocardic acids (e.g., C50) or mycolic acids (e.g., C90). Unless otherwise indicated, use of the term mycolic acids herein refers to any of these long chain α -substituted, β -hydroxy fatty acids. In particular, compounds according to Formula I exhibit antibiotic activity against corynebacteria, nocardiae, rhodococcus and mycobacteria. More particularly, these compounds are effective
15 against *Mycobacterium tuberculosis*, drug-resistant *M. tuberculosis*, *M. avium intracellulare*, *M. leprae*, or *M. paratuberculosis*.

Preferred compounds according to Formula I will have substantial antibiotic activity against susceptible organisms (see, e.g., Table 1). Antibiotic effectiveness of compounds according to Formula I may be determined as described
20 below or by use of assays described in U.S. Patent No. 5,614,551, which is incorporated herein by reference. In particular, U.S. Patent 5,614,551 describes an *in vitro* therapeutic index based on comparison of the concentration which inhibits growth of normal fibroblasts to the minimal inhibitory concentration for a compound, and preferred compounds will have an *in vitro* therapeutic index of at least 2, more preferably at least
25 5, and most preferably at least 10.

Novel drug therapy, using compounds of this invention which are effective against multiply drug resistant tuberculosis, will aid in treating both patients presently suffering from active disease, and the millions of potential patients who harbor quiescent disease which may become active as a result of immunosuppression

or other systemic disease. These drugs will also be useful against the "atypical mycobacteria" such as *M. avium-intracellulare*, a common AIDS pathogen, and other species that are commonly drug resistant. Given the biochemical similarity between *M. tuberculosis* and *M. leprae*, these drugs may be expected to be useful in the treatment of leprosy (Hansen's disease). Potential use in livestock or other veterinary applications include treatment of infections by *Mycobacterium paratuberculosis*, also known as Johne's bacillus, an organism that produces a chronic enteritis in ruminants (e.g., cattle and sheep) which is invariably fatal, and *Rhodococcus*, another organism which produces mycolic acids as well as potentially fatal respiratory infections in horses and immunocompromised patients. Treatment of human patients infected with *M. paratuberculosis* is also within the contemplation of this invention.

Treatment according to this invention involves administering the compound of Formula I to the subject of treatment. Pharmaceutical compositions containing any of the compounds of this invention may be administered by parenteral (subcutaneously, intramuscularly, intravenously, intraperitoneally, intrapleurally, intravesicularly or intrathecally), topical, oral, rectal, or nasal or inhalation route, as necessitated by choice of drug, pharmaceutical carrier, and disease.

Therapeutic compounds according to this invention are preferably formulated in pharmaceutical compositions containing the compound and a pharmaceutically acceptable carrier. The concentrations of the active agent in pharmaceutically acceptable carriers will depend on solubilities. The dose used in a particular formulation or application will be determined by the requirements of the particular type of disease and the constraints imposed by the characteristics and capacities of the carrier materials. The pharmaceutical composition may contain other components so long as the other components do not reduce the effectiveness of the compound according to this invention so much that the therapy is negated. Pharmaceutically acceptable carriers are well known, and one skilled in the pharmaceutical art can easily select carriers suitable for particular routes of

administration (see, e.g., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, PA, 1985).

Dose and duration of therapy will depend on a variety of factors, including the therapeutic index of the drugs, disease type, patient age, patient weight, and tolerance of toxicity. Dose will generally be chosen to achieve serum concentrations from about 1 ng to about 100 $\mu\text{g/ml}$, typically 0.1 $\mu\text{g/ml}$ to 10 $\mu\text{g/ml}$. Preferably, initial dose levels will be selected based on their ability to achieve ambient concentrations shown to be effective in *in vitro* models and *in vivo* models and in clinical trials, up to maximum tolerated levels. The dose of a particular drug and duration of therapy for a particular patient can be determined by the skilled clinician using standard pharmacological approaches in view of the above factors. The response to treatment may be monitored by analysis of blood or body fluid levels of the compound according to this invention, measurement of activity of the compound or its levels in relevant tissues or monitoring disease state in the patient. The skilled clinician will adjust the dose and duration of therapy based on the response to treatment revealed by these measurements.

Typically, the compositions described above will be combined or used together or in coordination with one or more other therapeutic substances, e.g., other drugs presently used in treating tuberculosis. The compound of Formula I, or a synergistic combination of inhibitors, will of course be administered at a level (based on dose and duration of therapy) below the level that would kill the animal being treated. Preferably, administration will be at a level that will not irreversibly injure vital organs, or will not lead to a permanent reduction in liver function, kidney function, cardiopulmonary function, gastrointestinal function, genitourinary function, integumentary function, musculoskeletal function, or neurologic function. On the other hand, administration of inhibitors at a level that kills some cells which will subsequently be regenerated (e.g., endometrial cells) is not necessarily excluded.

EXAMPLES

In order to facilitate a more complete understanding of the invention, a number of Examples are provided below. However, the scope of the invention is not limited to specific embodiments disclosed in these Examples, which are for purposes of illustration only.

Example 1. Synthesis of substituted sulfonylamides

To illustrate the synthetic method shown in Scheme 1, a 25 g synthesis of III-50 was carried out.

10 **Methyl n-Octylthioacetate.** Octyl bromide (50.21 g, 0.26 mole), methyl thioglycoate (22.35 ml, 26.53 g, 0.25 mole) and potassium carbonate (34.5 g, 0.25 mole) were charged into a 1 L round-bottomed flask. To this mixture was added 350 ml of acetone and the suspension was refluxed for 48 h. After cooling to room temperature, the reaction mixture was filtered with the aid of acetone. The filtrate was evaporated under
15 reduced pressure and the residue thus obtained was purified by distillation under reduced pressure. The fraction distilling at 118-121 °C/3.8 mm Hg was collected. Yield: 48 g; 88%

IR (neat): 2924, 1736, 1435, 1278, 1133, 1011 cm⁻¹.

¹H NMR (CDCl₃): δ 3.64 (s, 3H), 3.13 (s, 2H), 2.52 (t, 2H, J= 7.2 Hz), 1.47
20 (m, 2H), 1.2-1.4 (m, 10H), 0.88 (t, 3H, J= 6.8 Hz).

¹³C, NMR: (CDCl₃): δ 13.9, 22.5, 28.6, 29.0, 29.02, 31.6, 32.5, 33.3, 52.1, 171.0.

Methyl n-Octanesulphonylacetate. A 3 L three-necked round-bottomed flask (fitted with a mechanical stirrer) was charged sequentially with ammonium heptamolybdate tetrahydrate (56 g, 0.045 mole), methyl n-octylthioacetate (40 g, 0.183 mole) and 1.5
25 L absolute alcohol (Schultz, HS, Freyermuth, HB., and Buc, SR., *J. Org. Chem.* **28**:1140-1142, 1963). The vigorously stirred solution was cooled to 0 °C and to this cooled solution was added 104 ml of 30% hydrogen peroxide solution (0.732 mole) over a period of 1 h. The reaction mixture was allowed to warm to room temperature

over a period of 2 h and then stirred for another 24 h when thin layer chromatography on silica showed complete disappearance of the starting material. The reaction mixture was filtered and the filtrate was evaporated under reduced pressure. The residue thus obtained was dissolved in ethyl acetate (1 L) and washed with water (100 ml x 2), brine (100 ml). The organic layer was dried over anhydrous MgSO_4 and, after filtration of the solvents, was evaporated under reduced pressure to obtain the sulfone as a waxy solid, 36 g, 78%. The produce appeared to be pure (>95%) by NMR and was submitted to the next reaction without further purification.

10 **IR (neat):** 2919, 2848, 1743, 1460, 1437, 1329, 1278, 1218, 1137, 1108, 1010, 912, 723 cm^{-1} .

^1H NMR (CDCl_3): δ 3.96 (s, 2H), 3.79 (s, 3H), 3.21 (t, 3H, $J = 8\text{ Hz}$), 1.80 (m, 2H), 1.20-1.45 (m, 10H), 0.81 (t, 3H, $J = 6.7\text{ Hz}$).

^{13}C NMR (CDCl_3): δ 13.9, 21.7, 22.4, 28.2, 28.7, 28.8, 31.5, 53.1, 53.4, 57.0, 163.4.

15 **n-Octanesulphonylacetamide.** A solution of methyl n-octanesulphonylacacetate (35 g, 0.143 mole) in 350 ml of anhydrous methanol was stirred magnetically at room temperature. To this solution was added 24 ml of aqueous ammonium hydroxide (27%, 6.48 g, 0.185 mole) in drops over a period of 30 minutes. The solution was stirred for 24 h and the white precipitate formed was filtered. The solid was recrystallized from
20 hot ethyl acetate to obtain the required acetamide III-50 as a crystalline solid, mp. 140-142 $^{\circ}\text{C}$, 33g, 97.6%.

IR (neat): 3386, 2920, 1659, 1420, 1315, 1286, 1129, cm^{-1} .

25 **^1H NMR (CDCl_3):** δ 6.56 (br s, 1H, NH), 5.69 (br s, 1H, NH) 3.86 (s, 2H), 3.14 (app 1:1:1 triplet, $J_{\text{app}} = 8.0\text{ Hz}$, 2H), 2.16 (m, 2H), 1.86 (m, 2H), 1.1-1.3 (m, 12H), 0.86 (t, $J = 7.1\text{ Hz}$, 3H).


^{13}C NMR (CD_3COCD_3): δ 14.7, 22.9, 23.6, 29.4, 30.1, 30.15, 30.8, 32.8, 54.0, 164.8

HRMS calculated for $\text{C}_{10}\text{H}_{25}\text{N}_2\text{O}_3\text{S}$ ($\text{M}+\text{NH}_4^+$) 253.1586, found 253.1587.

Example 2. Synthesis of substituted sulfonylamides

To illustrate the synthetic methods shown in Scheme 2 below, the following syntheses of compounds I-31 and I-89 were carried out.

Scheme 2



- 5 **(*E/Z*)-3-decylsulfanyl-acrylic acid methyl ester (1):** Triethylamine (0.5 mL, 3.6 mmol) was added dropwise to a solution of decanethiol (2.5 mL, 12.1 mmol) and methyl propiolate (4.3 mL, 48.3 mmol) in dichloromethane (20 mL). The solution was stirred at room temperature for 20 min under argon, after which it was diluted with water (60 mL) and extracted with dichloromethane (3 x 60 mL). The organic layer was
- 10 washed with water (2 x 60 mL), brine (60 mL), dried over anhydrous magnesium sulfate, and concentrated *in vacuo*. Purification by flash chromatography (silica gel, ethyl acetate:hexanes 1:49) yielded a clear oil (3.06 g, 98%) as a mixture of isomers (*E:Z*, 6:1), which was carried through to the next step.
- 15 ***E* isomer:** IR (neat, cm^{-1}): 2920, 2848, 1712, 1580, 1460, 1430, 1300, 1255, 1215, 1160, 1041, 1015, 945, 827, 715, 697.
 ^1H NMR (CDCl_3) δ 0.88 (*t*, 3H, $J = 6.7$ Hz), 1.26 (*bs*, 12H), 1.40 (*m*, 2H), 1.67 (*quint*, 2H, $J = 7.3$ Hz), 2.78 (*t*, 2H, $J = 7.4$ Hz), 3.72 (*s*, 3H), 5.74 (*d*, 1H, $J = 15.2$ Hz), 7.70 (*d*, 1H, $J = 15.2$ Hz).
- 20 ***Z* isomer:** IR (neat, cm^{-1}): 2920, 2848, 1712, 1580, 1460, 1430, 1300, 1255, 1215, 1160, 1041, 1015, 945, 827, 715, 697.

¹H NMR (CDCl₃) δ 0.88 (t, 3H, J = 6.7 Hz), 0.9-1.9 (app. s & several m, 16H), 2.77 (t, 2H, J = 7.4 Hz), 3.75 (s, 3H), 5.85 (d, 1H, J = 10.2 Hz), 7.10 (d, 1H, J = 10.2 Hz).

(E)-3-decylsulfanyl-acrylic acid methyl ester (2): Compound 1 (0.50 g, 1.93 mmol), was dissolved in methanol (8.0 mL), and cooled to 0 °C, added to a solution of oxone (1.84 g, 6.0 mmol) in water (8.0 mL, 49.5%) at 0 °C, and stirred for 4 h at room temperature. The reaction was then diluted with water (60 mL), and extracted with chloroform (3 x 60 mL). The organic extract was washed with water (2 x 60 mL), brine (60 mL), dried over anhydrous magnesium sulfate, and concentrated *in vacuo*.
Purification by flash chromatography (silica gel, ethyl acetate:hexanes 1:1) yielded white crystals (0.53g, 93%) as a mixture of isomers (*E:Z*, 6:1). Purification by preparative TLC (ethyl acetate:hexanes 1:1) yielded compound 2 as white crystals.

mp 62-63 °C; IR (CH₂Cl₂, cm⁻¹): 3060, 2950, 2920, 2845, 1730, 1470, 1435, 1280, 1230, 1165, 1130, 995, 965, 815, 765, 690, 573.

¹H NMR (CDCl₃) δ 0.88 (t, 3H, J = 6.7 Hz), 1.27 (bs, 12H), 1.43 (m, 2H), 1.80 (sym. m, 2H), 3.05 (sym. m, 2H), 3.86 (s, 3H), 6.88 (d, 1H, J = 15.3 Hz), 7.34 (d, 1H, J = 15.3 Hz). Calculated for C₁₄H₂₆O₄S: C, 57.90, H, 9.02, S, 11.04. Found: C, 57.94, H, 9.12, S, 11.16.

20

Propiolamide (3): Methyl propiolate (2.0 mL, 22.5 mmol) was added to liquid ammonia at -78 °C and stirred for 2 hours. Evaporation at room temperature yielded compound 3 as white crystals (1.46 g, 94%).

mp 58-61°C (lit.[5], 61-62°C); ¹H NMR (D₂O) δ 3.50 (s, 1H), 4.43 (bs, 2H).

(E/Z)-3-decylsulfanyl-propionamide (4): Triethylamine (0.5 mL, 3.6 mmol) was added dropwise to a solution of decanethiol (1.36 mL, 6.56 mmol) and propiolamide (1.80 g, 26.1 mmol) in dichloromethane (20 mL). The solution was stirred

at room temperature for 20 min under argon, after which it was diluted with water (60 mL) and extracted with dichloromethane (3 x 60 mL). The organic layer was washed with water (2 x 60 mL), brine (60 mL), dried over anhydrous magnesium sulfate, and concentrated *in vacuo*. Purification by flash chromatography (silica gel, ethyl acetate:hexanes 3:1) yielded compound 4 as white crystals (1.45g, 91%) as a mixture of isomers (*E:Z*, 1:10), which was carried through to the next step.

Mixture of isomers: mp 65-70°C; IR (CH₂Cl₂, cm⁻¹): 3400, 3330, 3280, 3200, 2920, 2840, 1725, 1650, 1570, 1460, 1400, 1300, 1190, 770.

10 *E-isomer:* ¹H NMR (CDCl₃) δ 0.89 (*t*, 3H, *J* = 6.8 Hz), 1.27 (*bs*, 12H), 1.41 (*m*, 2H), 1.70 (*m*, 2H), 2.74 (*t*, 2H, *J* = 7.5 Hz), 5.4 (*bs*, 2H, NH₂), 5.82 (*d*, 1H, *J* = 10.1 Hz), 6.95 (*d*, 1H, *J* = 10.1 Hz).

(*E*)-3-decylsulfonyl-propionamide (5): Compound 4 (0.50 g, 2.05 mmol) was dissolved in methanol (8.0 mL), cooled to 0 °C and added to a solution of oxone (1.84 g, 6.0 mmol) in water (8.0 mL, 49.5%) at 0 °C and stirred for 4 h at room temperature. The solution was then diluted with water (60 mL), and extracted with chloroform (3 x 60 mL). The organic extract was washed with water (2 x 60 mL), brine (60 mL), dried over magnesium sulfate, and concentrated *in vacuo*. Purification by flash chromatography (silica gel, ethyl acetate:hexanes 17:3) yielded compound 5 as white crystals (0.52 g, 92%) as a mixture of isomers (*E:Z*, 6:1). Purification by preparative TLC (ethyl acetate:hexanes 3:1) yielded compound 5 as white crystals.

mp 148-149 °C; IR (CHCl₃, cm⁻¹): 3390, 3140, 3060, 2920, 2840, 1700, 1615, 1455, 1395, 1320, 1130, 960, 814.

¹H NMR (CDCl₃) δ 0.089 (*t*, 3H, *J* = 6.8 Hz), 1.27 (*bs*, 12H), 1.43 (*m*, 2H), 1.80 (*m*, 2H), 3.05 (*sym. m*, 2H), 5.67 (*bs*, 1H, NH), 5.85 (*bs*, 1H, NH), 6.94 (*d*, 1H, *J* = 14.8 Hz), 7.35 (*d*, 1H, *J* = 14.8 Hz).

Calculated for $C_{13}H_{25}NO_3S$: C, 56.70, H, 9.15, N 5.09, S 11.64. Found: C, 56.78, H, 9.16, N, 5.04, S, 11.76.

Example 3. *In vitro* activity of sulfones and sulfoxides against mycobacteria

5 Various compounds were tested as described in U.S. Patent No. 5,614,551 to determine minimum inhibitory concentrations (MIC) of the compounds against drug-resistant MTB (*M. tuberculosis* strain H37Rv), *M. avium-intracellulare*, and *M. bovis* BCG. The results are shown in Table 1. Dose-response curves for the compound, designated III-50, have demonstrated an MIC of 6.5 μ g/ml against the
10 virulent strain of *M. tuberculosis*, H37Rv, and 6.25 μ g/ml for *M. bovis* BCG. Dose-response curves for the compound designated S-I-73 have demonstrated an MIC of 3.12 μ g/ml against MTB and 12.50 μ g/ml against *M. avium-intracellulare*. Tests using a compound according to Formula I having $R = C_8H_{17}$, $Z = NH$ and $Y = -CO-O-CH_3$ against *M. tuberculosis*, H37Rv, demonstrated an MIC of 12.5 μ g/ml.

15

Example 4. III-50 inhibits mycolic acid synthesis via a target different from isoniazid.

In a series of metabolic labeling experiments, the activity of a number of lipid metabolic pathways were studied in the presence and absence of III-50 using
20 two-dimensional thin-layer chromatography and phosphorimage quantification. The TLC plates were spotted with mycobacterial acid methanolysates and developed in the first direction with petroleum ether (bp 60-80°C):acetone (95:5, v/v, 3 times) and in the second direction with toluene:acetone (97:3), v/v, once). Abbreviations in Figure 2 are: Ori = origin; A = α -mycolate; B = ketomycolate; and C = ω -mycolate. The left hand
25 panel shows acid methanolysates from *M. avium-intracellulare* control cultures. The right hand panel shows acid methanolysates from *M. avium-intracellulare* cultures treated with 12.5 μ g/ml n-octanesulphonylacetamide (III-50).


While minor qualitative and quantitative alterations of various lipid species were identified, the most profound effect was noted upon mycolic acid

synthesis. Reproductions of the phosphorimages (Figure 2) demonstrate that in the presence of III-50 mycolic acids are undetectable in *M. avium-intracellulare* and significantly reduced in *M. bovis* BCG. More importantly, while inhibition of mycolic acid synthesis is thought to be the mechanism of action of isoniazid (Takayama, et al., 1972 & 1975; Quemard, et al., 1993 & 1996; and Baldock, et al.), III-50 inhibits both the growth of *M. avium-intracellulare*, which is routinely isoniazid resistant (>2.5 µg/ml), and also an isoniazid (INH) resistant *M. tuberculosis* (>0.4 µg/ml). Thus, although both isoniazid and III-50 inhibit mycolic acid synthesis, the enzymatic target of III-50 within the mycolic acid synthetic pathway appears to differ from that of isoniazid.

Inhibition of mycolic acid synthesis leads to the disruption of the cell wall in mycobacteria. Figure 3 shows electron micrographs which depict cell division of *M. bovis* BCG in the presence (Panel A) and absence (Panel B) of III-50. Note, in the control, the well-developed cell wall and septum as the bacterium divides. In contrast, in the presence of III-50, there is disruption of the cell wall likely as a result of inhibition of mycolic acid synthesis.

Preparation of a combinatorial library of sulfonamide compounds

This invention contemplates combinatorial libraries that contains compounds of Formula II:




Formula II

where the various symbols have the same meaning as in Formula I, except that one or both of the vinyl hydrogens may be independently replaced by a group selected from alkyl, acyl, aryl, aralkyl, halogen; substituted or unsubstituted thiol; unsubstituted or substituted amino; hydroxy, and OR' wherein R' is selected from the group consisting of hydrogen, alkyl, acyl, aryl aralkyl, unsubstituted or substituted amino; substituted or unsubstituted thiol; and halogen; and a linear or cyclic carbon chain optionally interrupted with one or more heteroatom, and

- optionally substituted with one or more =O, or =S depending on the choice of electrophile in Scheme 3, or both vinyl hydrogens are replaced by a linear carbon chain to form a cyclic carbon moiety optionally interrupted with one or more heteroatoms, and optionally substituted with one or more =O, or =S. Typically,
- 5 the groups substituted at the vinyl positions will have from 1 to 20 carbons in aggregate, and the heteroatoms will generally be selected from B, N, O, P, and S, more usually N, O, and S. The combinatorial library of this invention may also include derivatives of Formula II produced by electrophilic or free radical addition at the double bond between the acyl group and the sulfur atom.
- 10 Using the method of synthesis of vinyl sulfones 2 and 5, it is possible to prepare collections of screenable compounds by combinatorial methods. This is illustrated in Scheme 3.

Scheme 3



- 15 The acidity of the acetylenic hydrogen in methyl propiolate or propionamide 3 allows its ready removal by base and reaction with electrophiles R_1-X (halides, aldehydes, ketones, esters, etc.) to give **6**. Thiols $\text{R}-\text{SH}$ can be added to **6** in the presence of mild base (i.e., potassium or sodium carbonate, triethylamine, etc.)

to give trans- or trans/cis-mixtures of variously substituted sulfanylacylic esters or amides. Oxidation selectively to sulfoxides **8** or sulfones **9** can be carried out. Cis- and trans-isomers can be separated by crystallization or chromatographically. By varying R_1 -X and R-SH or the identity of the ester or
5 amide, a combinatorial library can be prepared.

The use of combinatorial libraries of diverse chemical compounds in drug discovery is well known (Moos, et al. (1993), Ann. Rep. Med Chem., vol. 28, chap. 33, pp. 315-324, Academic Press; Gordon, et al. (1994), J. Med. Chem., 37:1385-1401). Methods of screening such libraries are described, e.g., in
10 Gordon, et al., and references cited therein. Combinatorial libraries comprising compounds according to this invention, including vinyl sulfones or sulfoxides prepared as described in Scheme 3, may be screened for biological activity by any suitable screening procedure.

In an exemplary screening procedure, members of a library could
15 be tested for ability to inhibit growth of *Toxoplasma gondii* using a test system described in U.S. Patent 5,614,551, incorporated herein by reference. For example, toxicity, fibroblast lysis and growth of *T. gondii* can be monitored in 24 well tissue culture plates containing human foreskin fibroblasts. Seven serial dilutions of three members of the library can be compared to three control wells
20 in each 24 well plate. Using multiple plates, large numbers of compounds from the library can be screened for their effect on *T. gondii*. Alternatively, using fewer dilutions, more compounds may be tested in each plate. Similar screening may be performed using *Mycobacterium* sp. in test systems described in U.S. Patent 5,614,551 to identify compounds in a combinatorial library which affect
25 mycobacteria. Other test systems suitable for screening compounds in combinatorial libraries according to this invention for their effects on other

pathogens and/or neoplastic cells are readily available to the skilled worker, in view of the teachings herein.

Alternatively, the library may be screened for inhibitory effect on an enzyme such as fatty acid synthase (FAS). Suitable assay procedures are
5 described in U.S. Patent No. 5,759,837, incorporated herein by reference, and multi-well plates may be used to perform simultaneous FAS assays in the presence and absence of numerous members of the library to compare the inhibitory effect of various member compounds. In another alternative, binding
affinity of member compounds for a particular receptor may be compared. Using
10 a suitable assay that measures progress of a biological process, the skilled worker can readily design a suitable screening procedure to screen the combinatorial library of this invention for biological effect.

For purposes of clarity of understanding, the foregoing invention has been described in some detail by way of illustration and example in conjunction with
15 specific embodiments, although other aspects, advantages and modifications will be apparent to those skilled in the art to which the invention pertains. The foregoing description and examples are intended to illustrate, but not limit the scope of the invention. Modifications of the above-described modes for carrying out the invention that are apparent to persons of skill in medicine, immunology, infectious diseases,
20 pharmacology, and/or related fields are intended to be within the scope of the invention, which is limited only by the appended claims.

All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference
25 to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.







Table 1. *In Vitro* Activities of Sulfones and Sulfoxides against *M. tuberculosis*

Compound #		MIC (ug/ml)		
		MTB	BCG	MAI
SI-73		3.12		12.50
HHI-50		6.25	6.25	25.00
SI-46		6.25		
SI-52		6.25		
HHI-206a		6.25		
HHI-206b		12.50		
SI-45		12.50	25.00	
HHI-302		>25		
SI-48		>25		
HHI-56		>25	>25	
DI-59		>25		

MTB = *M. tuberculosis* (strain H37Rv)BCG = *M. bovis*MAI = *M. avium-intracellulare*

See legend in Appendix for Methods.

Table 1 (continued). *In Vitro* Activities of Sulfones and Sulfoxides against *M. tuberculosis*

Compound #		MIC (ug/ml)		
		MTB	BCG	MAI
JRG-I-89-2		12.5		
JRG-I-89-1		25		
JRG-I-89-3 (mixture of cis and trans)		6.25		
JRG-I-31		12.5		
JRG-I-31 (mixture of cis and trans)		50		
JRG-I-43		50		

MTB = *M. tuberculosis* (strain H37Rv)

BCG = *M. bovis*


MAI = *M. avium-intracellulare*

Claims:

1. A compound having the formula:
$$R-SO_n-Z-CO-Y,$$

wherein:
 - 5 R is an alkyl group having 6-20 carbons;
Z is a radical selected from the group consisting of $-CH_2-$, $-O-$, $-NH-$, two of these radicals coupled together, and $-CH_2=CH_2-$;
Y is selected from $-NH_2$, $O-CH_2-C_6H_5$, $-CO-CO-O-CH_3$, and $O-CH_3$; and
n is 1 or 2.
- 10 2. The compound according to claim 1, wherein R is a branched alkyl group.
3. The compound according to claim 1, wherein R is a linear alkyl group interrupted by an aromatic ring.
- 15 4 A method of inhibiting growth of a microbial cell comprising treating the cell with a compound according to claim 1, wherein the microbial cell is a cell which synthesizes α -substituted, β -hydroxy fatty acids.
 - 20 5. The method of claim 4, wherein the α -substituted, β -hydroxy fatty acid is selected from the group consisting of corynemycolic acid, nocardic acid, and mycolic acid.
 6. The method of claim 4, wherein the microbial cell is selected from the group consisting of corynebacteria, nocardiae, rhodococcus, and mycobacteria.
 - 25 7. The method of claim 6, wherein the mycobacterial cell is *Mycobacterium tuberculosis*, drug-resistant *M. tuberculosis*, *M. avium intracellulare*, *M. leprae*, or *M. paratuberculosis*.
 8. A method for treating a mycobacterial infection by administering to an animal a pharmaceutical composition containing a compound according to claim 1.

9. A combinatorial library comprising a plurality of different members, each member selected from the group consisting of compounds having the formula:



5 wherein:

R is an alkyl group having 6-20 carbons;

R_1 and R_2 are each independently selected from the group consisting of hydrogen; alkyl, acyl, aryl, aralkyl, halogen; substituted or unsubstituted thiol; unsubstituted or substituted amino; hydroxy, and OR' wherein R' is selected
10 from the group consisting of hydrogen, alkyl, acyl, aryl aralkyl, unsubstituted or substituted amino; substituted or unsubstituted thiol; and halogen; and a linear or cyclic carbon chain optionally interrupted with one or more heteroatom, and optionally substituted with one or more =O, or =S, or R_1 and R_2 together form a cyclic carbon chain optionally interrupted with one or more heteroatom, and
15 optionally substituted with one or more =O, or =S;

Y is selected from $-NH_2$, $O-CH_2-C_6H_5$, $-CO-CO-O-CH_3$, and $O-CH_3$; and

n is 1 or 2.

10. A method of identifying an effector of a biological process comprising carrying out the biological process in the presence of a plurality of
20 compounds selected from the library of claim 9 and comparing the results of the process to the results of carrying out the process in the absence of said compounds.

11. A method of identifying a growth inhibitor effective against growth of a predetermined microbe comprising culturing the microbe in the presence and
25 absence of each of a plurality of member compounds selected from the library of claim 9 and identifying the member compounds in whose presence growth of the microbe is least.

1/4




FIG. 1

2/4




FIG. 2

3/4



FIG. 3A

4/4



FIG. 3B

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 311/06, 317/44, A61K 31/18, 31/22	A3	(11) International Publication Number: WO 99/10321 (43) International Publication Date: 4 March 1999 (04.03.99)
(21) International Application Number: PCT/US98/17830 (22) International Filing Date: 28 August 1998 (28.08.98) (30) Priority Data: 60/056,272 29 August 1997 (29.08.97) US (71) Applicant (for all designated States except US): THE JOHNS HOPKINS UNIVERSITY [US/US]; 720 Rutland Avenue, Baltimore, MD 21205 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): TOWNSEND, Craig, A. [US/US]; 116 Midhurst Road, Baltimore, MD 21212 (US). DICK, James, D. [US/US]; 15744 Dover Road, Upperco, MD 21155 (US). PASTERNAK, Gary, R. [US/US]; 311 Edgevale Road, Baltimore, MD 21212 (US). KUHAJDA, Francis, P. [US/US]; 1211 Broadway Road, Lutherville, MD 21209 (US). PARRISH, Nicole, M. [US/US]; Apartment 201, 3185 Pine Orchard Lane, Ellicott City, MD 21042 (US). (74) Agents: POSORSKE, Laurence, H. et al.; Baker & Botts, L.L.P., The Warner, 1299 Pennsylvania Avenue, N.W., Washington, DC 20004 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 30 September 1999 (30.09.99)
(54) Title: ANTIMICROBIAL COMPOUNDS (57) Abstract This invention provides methods for treating a mycobacterial infection by administering to an animal a pharmaceutical composition containing a compound having the formula $R-SO_n-Z-CO-Y$, where R is an alkyl group having 6-20 carbons; Z is a radical selected from $-CH_2-$, $-O-$, and $-NH-$, two of these radicals coupled together, or $-CH_2=CH_2-$; Y is $-NH_2$, $O-CH_2-C_6H_5$, $-CO-CO-O-CH_3$, or $O-CH_3$; and n is 1 or 2. It has been discovered that these compounds inhibit growth of microbial cells which synthesize α -substituted, β -hydroxy fatty acids, particularly corynemycolic acid, nocardic acid, and mycolic acid. These compounds may be used to inhibit growth of mycobacterial cells, such as <i>Mycobacterium tuberculosis</i> , drug-resistant <i>M. tuberculosis</i> , <i>M. avium intracellulare</i> , and <i>M. leprae</i> .		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/17830

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C311/06 C07C317/44 A61K31/18 A61K31/22

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 26 30 947 A (KAO SOAP CO LTD) 3 February 1977 (1977-02-03) see table 7, page 41, entry 2; page 46, entry 1; table 9, page 62, entry 2; page 64, entry 2; table, page 76, entries 3-4; page 80, entries 6-11; page 81, entry 1; table, page 87, entries 6-10; page 88, entries 1-2	1
A	see page 7, line 16 - page 8, line 5; examples 17-18	4
X	DE 27 29 685 A (KAO SOAP CO LTD) 5 January 1978 (1978-01-05) see page 8, lines 19, 22	1

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

15 March 1999

Date of mailing of the international search report

16. 08. 99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Amsterdam, L

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/17830

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	M. MIKOLAJCZYK ET AL: TETRAHEDRON, vol. 32, no. 8, 1976, pages 969-973, XP002096632 see table 4, compound 17 ---	1
X	C. GEORGES ET AL: J. CHEM. SOC., FARADAY TRANS. 1, vol. 84, no. 5, 1988, pages 1531-1542, XP002096633 see page 1541, compound 2d ---	1
X	US 5 530 113 A (W.J. CHRIST ET AL) 25 June 1996 (1996-06-25) see column 140, compound E2; column 141, compound E4; column 142, compound E6 ---	1
X	M.S.F. JIE ET AL: CHEM. PHYS. LIPIDS, vol. 61, no. 2, 1992, pages 139-147, XP002096634 see table I, compounds 2-3; table II, compounds 12-13 ---	1
X	DATABASE CHEMLIST [Online] AMERICAN CHEMICAL SOCIETY STN, accession no. 32917, XP002096636 see 2(dodecylsulphonyl)acetamide (EINECS) & OFFICIAL GAZETTE OF THE EUROPEAN COMMUNITIES, vol. ANNEX, 15 June 1990 (1990-06-15), ---	1
A	F.E. HARDY ET AL: J. CHEM. SOC. C, no. 17, 1969, pages 2334-2336, XP002096635 see page 2335, table, compound VIIa -& DATABASE CHEMABS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US STN, accession no. 72:6560, XP002096637 see abstract; RN 26139-12-8 ---	1
A	WO 95 19706 A (THE JOHNS HOPKINS UNIVERSITY) 27 July 1995 (1995-07-27) -----	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/ 17830

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 8
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See further information

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-8

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-8

Compounds of the general formula of claim 1 and their use in a method of inhibiting the growth of microbial cells which synthesize alpha-substituted beta-hydroxy fatty acids, such as corynebacteria, nocardiae, rhodococcus and mycobacteria

2. Claims: 9-10

A combinatorial library comprising a plurality of different compounds of the general formula of claim 9 and a method for identifying an effector of a biological process, such as a growth inhibitor effective against the growth of a predetermined microbe, using a plurality of compounds selected from this library

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/17830

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2630947 A	03-02-1977	JP 1039567 C	31-03-1981
		JP 52010219 A	26-01-1977
		JP 55030706 B	13-08-1980
		JP 1044220 C	30-04-1981
		JP 52010220 A	26-01-1977
		JP 55033713 B	02-09-1980
		CH 595338 A	15-02-1978
		GB 1557225 A	05-12-1979
DE 2729685 A	05-01-1978	JP 1048925 C	28-05-1981
		JP 53003520 A	13-01-1978
		JP 55046362 B	22-11-1980
		JP 1048928 C	28-05-1981
		JP 53003521 A	13-01-1978
		JP 55046365 B	22-11-1980
		CH 623724 A	30-06-1981
		FR 2356368 A	27-01-1978
		GB 1577229 A	22-10-1980
		NL 7706523 A	03-01-1978
		US 4123254 A	31-10-1978
US 5530113 A	25-06-1996	AU 660325 B	22-06-1995
		AU 2526692 A	22-04-1993
		CA 2079971 A	12-04-1993
		CN 1071668 A	05-05-1993
		EP 0536969 A	14-04-1993
		FI 924579 A	12-04-1993
		HU 64538 A	28-01-1994
		JP 5194470 A	03-08-1993
		NO 180491 B	20-01-1997
		NZ 244643 A	26-07-1995
		RU 2095366 C	10-11-1997
		US 5843918 A	01-12-1998
		US 5756718 A	26-05-1998
		US 5612476 A	18-03-1997
		ZA 9207751 A	10-06-1993
WO 9519706 A	27-07-1995	US 5614551 A	25-03-1997
		AU 703272 B	25-03-1999
		AU 1726895 A	08-08-1995
		CA 2181031 A	27-07-1995
		EP 0738107 A	23-10-1996
		JP 9508370 T	26-08-1997

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☒ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)